arXiv:1303.3436v1 [astro-ph.CO] 14 Mar 2013

Cosmology with Ricci dark energy

Sergio del Campo*,[1] Júlio.C. Fabris†,[2] Ramón Herrera‡,[1] and Winfried Zimdahl§[2]

[1] *Instituto de Física, Pontificia Universidad Católica de Valparaíso,*
Avenida Brasil 2950, Casilla 4059, Valparaíso, Chile

[2] *Universidade Federal do Espírito Santo, Departamento de Física*
Av. Fernando Ferrari, 514, Campus de Goiabeiras,
CEP 29075-910, Vitória, Espírito Santo, Brazil

(Dated: March 15, 2013)

Abstract

We assume the cosmological dark sector to consist of pressureless matter and holographic dark energy with a cutoff length proportional to the Ricci scale. The requirement of separate energy-momentum conservation of the components is shown to establish a relation between the matter fraction and the (necessarily time-dependent) equation-of-state parameter of the dark energy. Singling out intrinsically adiabatic pressure perturbations of the dark-energy component, the matter perturbations are found as linear combinations of the total energy-density perturbations of the cosmic medium and the relative (non-adiabatic) perturbations of the components. The resulting background dynamics is consistent with observations from supernovae of type Ia, baryonic acoustic oscillations and the differential age of old objects. For a physically interesting parameter space the perturbation quantities show a stable behavior. On smaller scales the model predicts oscillations in the dark sector which potentially should be testable by weak gravitational lensing and by their impact on the anisotropy spectrum of the cosmic microwave background.

---

* E-Mail: sdelcamp@ucv.cl
† E-mail: fabris@pq.cnpq.br
‡ E-mail: ramon.herrera@ucv.cl
§ E-mail: winfried.zimdahl@pq.cnpq.br

I. INTRODUCTION

Despite of the many efforts to consistently explain the results of the observations of supernovae of type Ia (SNIa) in [1], the physical nature of the dark sector of the Universe, assumed to consist of dark matter (DM) and dark energy (DE), remains largely mysterious. The favored cosmological model is the Λ-cold-dark-matter (ΛCDM) model which also serves as a reference for alternative approaches to the DE problem. *Grosso modo*, the ΛCDM model does well in fitting most observational data (see, e.g., the recent WMAP 9 results [2]). Nevertheless, there is an ongoing interest in alternative models within General Relativity (GR) itself and beyond it. Although no serious contender seems to be around at the present time, these efforts continue to make sense not only because of the notorious cosmological constant and coincidence problems (see, e.g. [3]) but also to test as many potential deviations from the "standard" description as possible in order to constrain additional parameter sets which are usually introduced in alternative approaches in order to quantify these deviations. Typically, these approaches "dynamize" the cosmological constant in terms of scalar fields or fluids with a generally time-dependent equation of state (EoS). Constraining a potential time dependence by the data, e.g., is then crucial for a comparison with the ΛCDM model. Holographic models represent a specific class of dynamic approaches to the DE problem. These models are characterized by a relation between an ultraviolet cutoff and an infrared cutoff [4–6]. Such relation guarantees that the energy in a given volume does not exceed the energy of a black hole of the same size. The infrared cutoff has to be a cosmological length scale. For the most obvious choice, the Hubble radius, only models in which dark matter and dark energy are interacting with each other also non-gravitationally, give rise to a suitable dynamics [7, 8]. Following [5], there has been a considerable number of investigations based on the future event horizon as cutoff scale [9]. However, all models with a cutoff at the future event horizon suffer from the serious drawback that they cannot describe a transition from decelerated to accelerated expansion. A future event horizon does not exist during the period of decelerated expansion. A further option that has received attention more recently and which will be the subject of the present paper is a model based on a cutoff length proportional to the Ricci scale. The role of a distance proportional to the Ricci scale as a causal connection scale for perturbations was noticed in [10]. As a cutoff length in DE models it was first used in [11]. Further investigations along this line have been carried out in [12]. In [13–15] observational constraints were obtained on the basis of which Ricci DE was compared with the ΛCDM model. The dynamics of perturbations was considered in [16–18]. A number of studies performed in [19–24] include interactions between DM and Ricci DE. A relation to quantum field theory has been claimed in [25].

In the present paper we reconsider the dynamics of a two-component system of pressureless DM and Ricci-type DE both in the homogeneous and isotropic background and on the perturbative level. While most dynamic DE scenarios start with an assumption for the EoS parameter for the DE, the starting point of holographic models is an expression for the DE energy density from which the EoS is then derived. Moreover, as was pointed out in [26], the mere definition of the holographic DE density generally implies an interaction with the DM component. Requiring this interaction to vanish imposes an additional condition on the dynamics. Non-interacting Ricci-type DE in particular is characterized by a simple relation between the matter fraction and the (time-dependent) EoS parameter. We shall confront the resulting background dynamics with recent SNIa data, results from baryonic acoustic oscillations (BAO) and from the history $H(z)$ of the Hubble parameter. Based on a gauge-invariant analysis, the matter perturbations are found as a combination of the total and the relative energy-density perturbations.

The structure of the paper is as follows. In Section II we recall basic relations for holographic models of DE. The resulting homogeneous and isotropic background dynamics is confronted with observational data in Section III. Section IV provides us with the general two-component dynamics of the cosmic medium. The first-order perturbation theory of the model is presented in sections V, VI and VII. On this basis, the final set of coupled equations for the non-adiabatic perturbation dynamics is found in section VIII. A summary of the paper is given in the final section IX.

II. BACKGROUND DYNAMICS FOR RICCI DARK ENERGY

We start by recalling basic features of holographic DE models in a homogeneous and isotropic background [26]. The cosmic medium is assumed to be describable by pressureless DM with energy density ρ_m and a holographic DE component with energy density ρ_H. In the spatially flat case Friedmann's equation is

$$3H^2 = 8\pi G(\rho_m + \rho_H) . \qquad (1)$$

In general, both components are not necessarily conserved separately but obey the balance equations

$$\dot{\rho}_m + 3H\rho_m = Q \ , \qquad \dot{\rho}_H + 3H(1+\omega)\rho_H = -Q \, , \tag{2}$$

such that the total energy $\rho = \rho_m + \rho_H$ is conserved. Here, $\omega \equiv \frac{p_H}{\rho_H} = \frac{p}{\rho_H}$ is the equation-of-state (EoS) parameter of the DE and p_H is the pressure associated with the holographic component. The acceleration equation can be written

$$\dot{H} = -\frac{3}{2}H^2\left(1 + \frac{\omega}{1+r}\right) \ , \quad \Rightarrow \quad \frac{d\ln H}{d\ln a} = -\frac{3}{2}\left(1 + \frac{\omega(a)}{1+r(a)}\right) \ , \tag{3}$$

where $r \equiv \frac{\rho_m}{\rho_H}$ is the ratio of the energy densities. The *total* effective EoS of the cosmic medium is

$$\frac{p}{\rho} = \frac{\omega}{1+r} \ . \tag{4}$$

According to the balance equations (2), the ratio r changes as

$$\dot{r} = 3Hr\ (1+r)\ \left[\frac{\omega}{1+r} + \frac{Q}{3H\rho_m}\right] \ . \tag{5}$$

Following [4, 5], we write the holographic energy density as

$$\rho_H = \frac{3\,c^2\,M_p^2}{L^2} \ . \tag{6}$$

The quantity L is the infrared (IR) cutoff scale and $M_p = 1/\sqrt{8\pi\,G}$ is the reduced Planck mass. The numerical constant c^2 determines the degree of saturation of the condition

$$L^3\,\rho_H \leq M_{Pl}^2\,L \, , \tag{7}$$

which is crucial for any holographic DE model. It states that the energy in a box of size L should not exceed the energy of a black hole of the same size [4].

Differentiation of the expression (6) and use of the energy balances (2) yields

$$\frac{Q}{\rho_H} = 2\frac{\dot{L}}{L} - 3H\,(1+\omega) \ . \tag{8}$$

In general, there is no reason for Q to vanish. Assuming $Q = 0$ provides us with a specific relationship between ω and the ratio of the rates $\frac{\dot{L}}{L}$ and H. Any non-vanishing Q will modify this relationship.

With Q from (8), the general dynamics (5) of the energy density ratio r becomes

$$\dot{r} = -3H\ (1+r)\ \left[1 + \frac{\omega}{1+r} - \frac{2}{3}\frac{\dot{L}}{HL}\right] \ . \tag{9}$$

The case without interaction is characterized by (cf. eq. (5))

$$Q = 0 \quad \Rightarrow \quad \dot{r} = r\left(2\frac{\dot{L}}{L} - 3H\right) = 3H\,r\,\omega \tag{10}$$

with a generally time-dependent ω. Different choices of the cutoff scale L give rise to different expressions for the total effective EoS parameter in (4) and to different relations between ω and r. Our interest in the present paper will be the Ricci-scale cutoff. The role of a distance proportional to the Ricci scale as a causal connection scale for perturbations was noticed in [10]. In [11] it was used for the first time as a DE cutoff scale. The Ricci scalar is $R = 6\left(2H^2 + \dot{H}\right)$. For the corresponding cutoff-scale one has $L^2 = 6/R$, i.e.,

$$\rho_H = 3\,c^2\,M_p^2\,\frac{R}{6} = \alpha\left(2H^2 + \dot{H}\right)\ , \tag{11}$$

where $\alpha = \frac{3c^2}{8\pi G}$. Upon using (3) we obtain

$$\rho_H = \frac{\alpha}{2}\,H^2\left(1 - 3\frac{\omega}{1+r}\right) \tag{12}$$

for the holographic DE density. Notice that the (not yet known) EoS parameter explicitly enters ρ_H. Use of Friedmann's equation provides us with

$$1 = \frac{c^2}{2}\left(1 + r - 3\omega\right) \quad \Rightarrow \quad \omega = \frac{1}{3}\left(1 + r\right) - \frac{2}{3c^2}\ . \tag{13}$$

Obviously, a constant value of ω necessarily implies a constant r and vice versa. The time derivatives of ω and r are related by $\dot{r} = 3\dot{\omega}$. The second relation in (13) can be used to express c^2 in terms of the present values (subindex 0) of ω and r:

$$\frac{2}{c^2} = 1 + r_0 - 3\omega_0\ , \quad \Rightarrow \quad r = r_0 + 3(\omega - \omega_0)\ . \tag{14}$$

The parameter c is related both to r_0 and ω_0. In a next step we differentiate ρ_H in (11) which yields

$$\dot{\rho}_H = \alpha\left(4H\dot{H} + \ddot{H}\right)\ . \tag{15}$$

With the help of (3) and the definition (11) with we derive [26]

$$\dot{\rho}_H + 3H\left(1 + \omega\right)\rho_H = -Q\ , \tag{16}$$

where

$$Q = -\frac{3H}{1+r}\left[r\omega - \frac{\dot{\omega}}{H}\right]\rho_H\ . \tag{17}$$

Relation (16) with (17), which implies that in the general case one has $Q \neq 0$, i.e., both dark components do interact with each other also non-gravitationally, is a direct consequence of the ansatz (11). The DE balance in Eq. (2) may then also be written as $\dot{\rho}_H + 3H\left(1+\omega_{eff}\right)\rho_H = 0$ with an effective EoS parameter

$$\omega_{eff} = \frac{1}{1+r}\left(\omega + \frac{\dot{\omega}}{H}\right) = \frac{\omega + \frac{\dot{\omega}}{H}}{1 + r_0 + 3\left(\omega - \omega_0\right)} \ . \tag{18}$$

The present ratio r_0 is related to the present matter fraction Ω_{m0} of the Universe by $r_0 = \frac{\Omega_{m0}}{1-\Omega_{m0}}$.

According to relation (17), a constant EoS parameter ω is compatible with $Q = 0$ only for $\omega = 0$, i.e., if ρ_H behaves as dust. If we admit $\dot{\omega} \neq 0$, however, there exists a non-trivial case $Q = 0$:

$$Q = 0 \quad \Rightarrow \quad r\omega = \frac{\dot{\omega}}{H} \quad \Rightarrow \quad r = \frac{d\ln\omega}{d\ln a} \ . \tag{19}$$

It is this configuration that we shall investigate in the present paper. Equation (19) together with the second relation of (14) is a differential equation for ω which has the solution

$$\omega = \omega_0 \frac{r_0 - 3\omega_0}{r_0 a^{-(r_0 - 3\omega_0)} - 3\omega_0} \ , \tag{20}$$

where we have normalized the present value of the cosmic scale factor a to $a_0 = 1$. There is no freedom left to choose the equation of state. It is fixed by the choice of ρ_H together with the requirement $Q = 0$. Notice that this is different from the more familiar procedure to deal with (non-holographic) DE, where one starts with an assumption for the EoS parameter and afterwards finds an expression for the DE density by integrating the corresponding balance equation. Here, the starting point is the energy density and the EoS parameter has to be derived.

Knowing the EoS parameter (20), it follows from (14) that

$$r = r_0 \frac{r_0 - 3\omega_0}{r_0 - 3\omega_0 a^{(r_0 - 3\omega_0)}} \ , \tag{21}$$

i.e., $r(a)$ is fixed as well. At high redshifts we have

$$\omega \to 0 \ , \qquad r \to r_0 - 3\omega_0 \ , \qquad\qquad (a \ll 1) \ . \tag{22}$$

The property that non-interacting Ricci-DE behaves as dust at high redshift was already pointed out in [11]. The values in the far-future limit are

$$\omega \to \omega_0 - \frac{1}{3}r_0 \ , \qquad r \to 0 \ , \qquad\qquad (a \gg 1) \ . \tag{23}$$

The limits in (22) imply that this model naturally reproduces an early matter-dominated era. For $r_0 \approx \frac{1}{3}$ and $\omega_0 \approx -1$, the ratio r approaches $r \approx \frac{10}{3}$ for $a \ll 1$. This value is only roughly ten times larger than the present value r_0. For the ΛCDM model the corresponding difference is about nine orders of magnitude. In this sense, the coincidence problem is considerably alleviated for the the present model. On the other hand, in the opposite limit $a \gg 1$ the ratio r approaches zero as for the ΛCDM model. Apparently, the far-future EoS can be of the phantom type for $\omega_0 - \frac{1}{3}r_0 < -1$.

The Hubble rate of our model turns out to be

$$\frac{H}{H_0} = a^{-3/2}\sqrt{\frac{3\omega_0 a^{(r_0-3\omega_0)} - r_0\left[1 + r_0 - 3\omega_0\right]}{3\omega_0 - r_0\left[1 + r_0 - 3\omega_0\right]}} \ . \tag{24}$$

For $a \ll 1$ we recover the Einstein - de Sitter behavior $H \propto a^{-3/2}$. The total effective EoS is

$$\frac{p}{\rho} = \frac{\omega}{1+r} = \omega_0\,\frac{r_0 - 3\omega_0}{r_0 a^{-(r_0-3\omega_0)}\left[1 + r_0 - 3\omega_0\right] - 3\omega_0} \ . \tag{25}$$

For the adiabatic sound speed of the DE component we find

$$\frac{\dot{p}_H}{\dot{\rho}_H} = \omega\left(1 - \frac{1}{3}\frac{r}{1+\omega}\right) \tag{26}$$

and the corresponding quantity of the total cosmic medium is

$$\frac{\dot{p}}{\dot{\rho}} = \frac{\dot{p}_H}{\dot{\rho}} = \omega\frac{1 - \frac{1}{3}\frac{r}{1+\omega}}{1 + \frac{r}{1+\omega}} \ . \tag{27}$$

With the solutions (20) and (21) all these quantities are explicitly known, i.e., the background dynamics is completely solved analytically. In the following section we perform an actualized confrontation of the background dynamics with recent observational data.

III. OBSERVATIONAL ANALYSIS

Our observational analysis of the background dynamics uses the following three tests: the differential age of old objects based on the $H(z)$ dependence as well as the data from SNIa and from BAO. A fourth test could potentially be added: the position of the first peak of the anisotropy spectrum of the cosmic microwave background radiation (CMB). However, the CMB test implies integration of the background equations until $z \sim 1.000$ which requires the introduction of the radiative component. But the inclusion of such radiative component considerably changes the structure of the equations and no analytic expression for $H(z)$ is available. Hence, we shall limit ourselves to the mentioned three tests for which a reliable estimation is possible.

Based on the evaluation of the age of old galaxies that have evolved passively, there are 13 observational data available for the differential age [27–31]. Recently, a new set of 21 data has been considered [32]. The basic relation is

$$H(z) = -\frac{1}{z}\frac{dz}{dt} \ . \tag{28}$$

The value of the Hubble parameter today can be added to these data, leading to 14 or 22 observational points, depending on the sample used.

The SNIa test is based on the distance modulus μ which is related to the luminosity distance D_L by

$$\mu = m - M = 5\log_{10} D_L \ , \qquad D_L = (1+z)\,\frac{c}{H_0}\int_0^z \frac{dz'}{\sqrt{H\left(z'\right)}} \ . \tag{29}$$

In this expression we have restored the velocity of light c. The quantities m and M denote the apparent and the absolute magnitudes, respectively.

Two decisions have to be taken for this test. The first one concerns the choice of the sample. There are many different SNIa data sets, obtained with different techniques. In some cases, these different samples may give very different results. The second point is the existence of two different calibration methods. One of them uses cosmological relations and takes into account SNIa with high z (Salt 2), the other one, using astrophysical methods, is suitable for small z (MLCS2k2) [33]. In some cases, the application of different calibrations can lead to different results also. All this makes the SNIa analysis very delicate. Here, we use the Union 2 sample [34], calibrated by the Salt 2 method.

Baryonic acoustic oscillations have their origin in oscillations in the photon-baryon plasma at the moment of the decoupling at about $z = 1.090$. They can be characterized by the distance scale [35],

$$\mathcal{A} = \frac{\sqrt{\Omega_{m0}}}{[H(z_b)]^{1/3}}\left[\frac{1}{z_b}\int_0^{z_b}\frac{dz}{H(z)}\right]^{2/3} . \tag{30}$$

We shall use the WiggleZ-data [36] $\mathcal{A} = 0.474 \pm 0.034,\ 0.442 \pm 0.020$ and 0.424 ± 0.021 for the redshifts $z_b = 0.44,\ 0.60$ and 0.73, respectively.

Generally, the key quantity of a statistical analysis is the χ^2 parameter

$$\chi^2(x^j) = \sum_i^n \frac{(f(x^j)^t_i - f(x^j)^o_i)^2}{\sigma_i}, \tag{31}$$



FIG. 1: One dimensional PDFs for ω_0 (upper panels) and r_0 (lower panels) using, from left to right, $H(z)$, $SNIa$, BAO and the combination of the three tests.

where $f(x^j)_i^t$ is the theoretical evaluation of a given observable, depending on x^j free parameters, $f(x^j)_i^o$ is the corresponding observational value with an error bar σ_i and n is the total number of observational data for the given test. In terms of the χ^2 parameter one defines the probability distribution function (PDF) by

$$P(x^j) = A\, e^{-\chi^2/2}, \tag{32}$$

where A is a normalization constant. The estimations for one or for two given parameters are obtained by integrating over the remaining ones. For a combination of all tests we use the total χ^2-value χ^2_T,

$$\chi^2_T = \chi^2_{H(z)} + \chi^2_{SN} + \chi^2_{BAO}. \tag{33}$$

Assuming a spatially flat universe, the three free parameters of the model are the density-ratio parameter $r_0 = \frac{\Omega_{mo}}{1-\Omega_{m0}}$, the EoS ω_0 and the reduced Hubble parameter h, defined by $H_0 = 100\, h\, \mathrm{km/s/Mpc}$.

In figure 1 we display the one-dimensional PDFs for each of the tests and for their combination. The results for the density parameter are different for each test. The combination of all tests leads to a value of $r_0 \sim 0.4$, corresponding to $\Omega_{m0} \sim 0.29$, roughly in agreement with the ΛCDM model. For the equation of state parameter we obtain



FIG. 2: Two-dimensional PDF for ω_0 and r_0 resulting from a combination of the three tests.

$\omega_0 \sim -1$, consistent with the ΛCDM model as well. According to the first relation of (14), the parameter c^2 turns out to be $c^2 \sim 0.46$. This value is coincides with the result in [11]. The two-dimensional PDFs at 1σ (68% of confidence level), 2σ (95% of confidence level) and 3σ (99% of confidence level) are shown in figure 2. The estimation for ω_0, based on a combination of the three tests at 2σ, is $\omega_0 = -0.987^{+0.083}_{-0.100}$, while for r_0 we find $r_0 = 0.406^{+0.073}_{-0.061}$.

IV. GENERAL TWO-COMPONENT DYNAMICS

To study the dynamics of inhomogeneities on the homogeneous and isotropic background of the previous sections we first consider the general description of the two-component system. It is characterized by a total energy-momentum tensor

$$T_{ik} = \rho u_i u_k + p h_{ik} \ , \qquad T^{ik}_{\ ;k} = 0, \tag{34}$$

where $h_{ik} = g_{ik} + u_i u_k$ and $g_{ik} u^i u^k = -1$. The quantity u^i denotes the total four-velocity of the cosmic substratum. Latin indices run from 0 to 3. The total T_{ik} splits into a matter component (subindex m) and a (holographic) DE component (subindex H),

$$T^{ik} = T^{ik}_m + T^{ik}_H, \tag{35}$$

with $(A = m, H)$

$$T^{ik}_A = \rho_A u^i_A u^k_A + p_A h^{ik}_A \ , \qquad h^{ik}_A = g^{ik} + u^i_A u^k_A \ . \tag{36}$$

For separately conserved fluids we have

$$T^{ik}_{m\ ;k} = 0, \qquad T^{ik}_{H\ ;k} = 0 \ . \tag{37}$$

Then, the separate energy conservation equations are

$$- u_{mi} T^{ik}_{m\ ;k} = \rho_{m,a} u^a_m + \Theta_m \rho_m = 0 \tag{38}$$

and

$$- u_{Hi} T^{ik}_{H\ ;k} = \rho_{H,a} u^a_H + \Theta_H \left(\rho_H + p_H\right) = 0 \ . \tag{39}$$

In general, each component has its own four-velocity, with $g_{ik} u^i_A u^k_A = -1$. The quantities Θ_A are defined as $\Theta_A = u^a_{A;a}$. For the homogeneous and isotropic background we assume $u^a_m = u^a_H = u^a$. Likewise, the momentum conservations are written as

$$h^a_{mi} T^{ik}_{m\ ;k} = \rho_m \dot{u}^a_m = 0 \tag{40}$$

and

$$h^a_{Hi} T^{ik}_{H\ ;k} = \left(\rho_H + p_H\right) \dot{u}^a_H + p_{H,i} h^{ai}_H = 0, \tag{41}$$

where $\dot{u}^a_A \equiv u^a_{A;b} u^b_A$.

Denoting first-order perturbations about the homogeneous and isotropic background by a hat symbol, the perturbed time components of the four-velocities are

$$\hat{u}_0 = \hat{u}^0 = \hat{u}^0_m = \hat{u}^0_H = \frac{1}{2}\hat{g}_{00} \ . \tag{42}$$

According to the perfect-fluid structure of both the total energy-momentum tensor (34) and the energy-momentum tensors of the components in (36), and with $u^a_m = u^a_H = u^a$ in the background, we have first-order energy-density perturbations $\hat{\rho} = \hat{\rho}_m + \hat{\rho}_H$, pressure perturbations $\hat{p} = \hat{p}_m + \hat{p}_H = \hat{p}_H$ and

$$\hat{T}^0_\alpha = \hat{T}^0_{m\alpha} + \hat{T}^0_{H\alpha} \quad \Rightarrow \quad \left(\rho + p\right) \hat{u}_\alpha = \rho_m \hat{u}_{m\alpha} + \left(\rho_H + p_H\right) \hat{u}_{H\alpha} \ . \tag{43}$$

In linear order the spatial components of the 4− accelerations are

$$\hat{\dot{u}}_\mu = \dot{\hat{u}}_\mu - \frac{1}{2}\hat{g}_{00,\mu}, \quad \hat{\dot{u}}_{(m)\mu} = \dot{\hat{u}}_{(m)\mu} - \frac{1}{2}\hat{g}_{00,\mu}, \quad \hat{\dot{u}}_{(H)\mu} = \dot{\hat{u}}_{(H)\mu} - \frac{1}{2}\hat{g}_{00,\mu} \ . \tag{44}$$

For the first-order pressure gradient terms we find (recall $p_H = p$)

$$\widehat{p_{,c}h^c_{H\mu}} = \hat{p}_{,\mu} + \dot{p}\hat{u}_{H\mu}, \qquad \widehat{p_{,c}h^c_\mu} = \hat{p}_{,\mu} + \dot{p}\hat{u}_\mu \ . \tag{45}$$

From the matter-momentum conservation (40) it follows that

$$\dot{u}^a_m = 0 \quad \Rightarrow \dot{\hat{u}}_{(m)\mu} - \frac{1}{2}\hat{g}_{00,\mu} = 0 \quad \Rightarrow \dot{\hat{u}}_{(m)\mu} = \frac{1}{2}\hat{g}_{00,\mu} \ . \tag{46}$$

According to (43) the differences between $\hat{u}_\alpha$ and the corresponding quantities of the components are

$$\hat{u}_\alpha - \hat{u}_{H\alpha} = \frac{\rho_m}{\rho + p}\left(\hat{u}_{m\alpha} - \hat{u}_{H\alpha}\right) \ , \quad \hat{u}_\alpha - \hat{u}_{m\alpha} = \frac{\rho_H + p_H}{\rho + p}\left(\hat{u}_{H\alpha} - \hat{u}_{m\alpha}\right) \ . \tag{47}$$

Restricting ourselves to scalar perturbations, the perturbed line element can be written

$$\mathrm{d}s^2 = -\left(1 + 2\phi\right)\mathrm{d}t^2 + 2a^2F_{,\alpha}\mathrm{d}t\mathrm{d}x^\alpha + a^2\left[\left(1 - 2\psi\right)\delta_{\alpha\beta} + 2E_{,\alpha\beta}\right]\mathrm{d}x^\alpha\mathrm{d}x^\beta \ . \tag{48}$$

Furthermore, we define the (three-) scalar quantities v, v_m and v_H by

$$a^2\hat{u}^\mu + a^2F_{,\mu} = \hat{u}_\mu \equiv v_{,\mu} \tag{49}$$

and

$$a^2\hat{u}^\mu_m + a^2F_{,\mu} = \hat{u}_{m\mu} \equiv v_{m,\mu} \ , \quad a^2\hat{u}^\mu_H + a^2F_{,\mu} = \hat{u}_{H\mu} \equiv v_{H,\mu} \ , \tag{50}$$

respectively.

From the definitions of Θ, Θ_m and Θ_H it follows that

$$\hat{\Theta} = \frac{1}{a^2}\left(\Delta v + \Delta\chi\right) - 3\dot{\psi} - 3H\phi \ , \tag{51}$$

where Δ is the three-dimensional Laplacian,

$$\chi \equiv a^2\left(\dot{E} - F\right) \tag{52}$$

and

$$\hat{\Theta}_m = \frac{1}{a^2}\left(\Delta v_m + \Delta\chi\right) - 3\dot{\psi} - 3H\phi \ , \quad \hat{\Theta}_H = \frac{1}{a^2}\left(\Delta v_H + \Delta\chi\right) - 3\dot{\psi} - 3H\phi \ . \tag{53}$$

V. CONSERVATION EQUATIONS IN LINEAR ORDER

At first order, the energy balances (38) and (39) are

$$\dot{\hat{\rho}}_m + \dot{\rho}_m \hat{u}^0 + \hat{\Theta}_m \rho_m + \Theta \hat{\rho}_m = 0 \tag{54}$$

and

$$\dot{\hat{\rho}}_H + \dot{\rho}_H \hat{u}^0 + \hat{\Theta}_H \left(\rho_H + p_H\right) + \Theta \left(\hat{\rho}_H + \hat{p}_H\right) = 0 \ , \tag{55}$$

respectively. The total first-order energy conservation takes the form

$$\dot{\hat{\rho}} + \dot{\rho} \hat{u}^0 + \hat{\Theta} \left(\rho + p\right) + \Theta \left(\hat{\rho} + \hat{p}\right) = 0 \ . \tag{56}$$

Comparing (54), (55) and (56) one finds

$$\hat{\Theta} = \frac{\rho_m}{\rho + p} \hat{\Theta}_m + \frac{\rho_H + p_H}{\rho + p} \hat{\Theta}_H \tag{57}$$

with $p = p_H$. The separate momentum conservation equations are given by (40) and (41). Additionally, we have the total momentum conservation

$$\left(\rho_m + \rho_H + p_H\right) \dot{u}^a + p_{H,i} h^{ai} = 0 \ . \tag{58}$$

For the total energy-density perturbations $\delta \equiv \frac{\hat{\rho}}{\rho}$ equation (56) yields

$$\dot{\delta} - \Theta \frac{p}{\rho} \delta - \frac{\dot{\rho}}{\rho} \phi + \hat{\Theta} \left(1 + \frac{p}{\rho}\right) + \Theta \frac{\hat{p}}{\rho} = 0 \ , \tag{59}$$

where we have used $\hat{u}^0 = -\phi$.

The momentum balance (58) for the cosmic medium as a whole together with the first relation of (44) and the second relation of (45) provides us with

$$\left(\rho + p\right) \left[\dot{v} + \phi\right] + \hat{p}^c = 0 \ , \tag{60}$$

where $\hat{p}^c = \hat{p} + \dot{p} v$.

In terms of the fractional perturbation $\delta_m \equiv \frac{\hat{\rho}_m}{\rho_m}$, the matter energy conservation (54) can be written as

$$\dot{\delta}_m - \frac{\dot{\rho}_m}{\rho_m} \phi + \hat{\Theta}_m = 0 \ . \tag{61}$$

The matter momentum balance (40) together with the second relation of (44) results in

$$\dot{v}_m + \phi = 0 \ . \tag{62}$$

With $\delta_H \equiv \frac{\hat{\rho}_H}{\rho_H}$ the energy conservation (55) for the DE component is

$$\dot{\delta}_H + \frac{\dot{\rho}_H}{\rho_H}\delta_H - \frac{\dot{\rho}_H}{\rho_H}\phi + \hat{\Theta}_H\left(1 + \frac{p_H}{\rho_H}\right) + \Theta\left(\delta_H + \frac{\hat{p}_H}{\rho_H}\right) = 0 \ . \tag{63}$$

In the following it will be convenient to introduce the quantity

$$D_H \equiv \frac{\hat{\rho}_H}{\rho_H + p_H} = \frac{\delta_H}{1+\omega} \ . \tag{64}$$

In terms of D_H eq. (63) then takes the form

$$\dot{D}_H - \Theta\frac{\dot{p}_H}{\dot{\rho}_H}D_H + \Theta\phi + \hat{\Theta}_H + \Theta\frac{\hat{p}_H}{\rho_H + p_H} = 0 \ . \tag{65}$$

The dark-energy momentum balance (41) together with the third relation of (44) and the first relation of (45) result in

$$\left(\rho_H + p_H\right)\left[\dot{v}_H + \phi\right] + \hat{p}^{c_H} = 0 \quad \Rightarrow \quad \dot{v}_H + \phi = -\frac{\hat{p}^{c_H}}{\rho_H + p_H} \equiv -P^{c_H} \ , \tag{66}$$

where $\hat{p}^{c_H} = \hat{p}_H + \dot{p}_H v_H = \hat{p} + \dot{p} v_H$.

Our final goal in this paper is to calculate the matter-density perturbations. To this purpose we shall solve the coupled system of total energy perturbations δ and relative energy-density perturbations $\delta_m - D_H$. In the following section we start by establishing the equation for the total energy-density perturbations.

VI. TOTAL ENERGY-DENSITY PERTURBATIONS

We consider equation (59) and introduce therein the gauge-invariant quantities

$$\delta^c = \delta + \frac{\dot{\rho}}{\rho}v \ , \quad \hat{\Theta}^c = \hat{\Theta} + \dot{\Theta}v \ . \tag{67}$$

Then, eq. (59) is rewritten as

$$\dot{\delta}^c - \Theta\frac{p}{\rho}\delta^c - \frac{\dot{\rho}}{\rho}\left(\dot{v} + \phi\right) + \frac{\Theta}{\rho}\hat{p}^c + \left(1 + \frac{p}{\rho}\right)\hat{\Theta}^c = 0 \ . \tag{68}$$

Combination of the energy conservation (68) and the momentum conservation (60) yields

$$\dot{\delta}^c - \Theta\frac{p}{\rho}\delta^c + \left(1 + \frac{p}{\rho}\right)\hat{\Theta}^c = 0 \ . \tag{69}$$

The perturbation $\hat{\Theta}$ has to be determined from the perturbed Raychaudhuri equation for Θ. Neglecting shear and vorticity, the Raychaudhuri equation is

$$\dot{\Theta} + \frac{1}{3}\Theta^2 - \dot{u}^a_{;a} + 4\pi G\left(\rho + 3p\right) = 0 \ . \tag{70}$$

In terms of the gauge-invariant variables one finds, at linear order,

$$\dot{\hat{\Theta}}^c + \frac{2}{3}\Theta\hat{\Theta}^c + 4\pi G\rho\delta^c - \dot{u}^a_{:a} = 0\ . \tag{71}$$

In a next step we have to differentiate (69) and to insert (71) into the resulting expression. The remaining $\hat{\Theta}^c$ terms can be eliminated by (69) again. Using also

$$\dot{u}^m_{;m} = -\frac{1}{a^2}\left(\frac{\Delta\hat{p} + \dot{p}\Delta v}{\rho + p}\right) = -\frac{1}{a^2}\frac{\Delta\hat{p}^c}{\rho + p}\ , \tag{72}$$

the equation for δ^c becomes

$$\ddot{\delta}^c + \left[2 - 6\frac{p}{\rho} + 3\frac{\dot{p}}{\dot{\rho}}\right]H\dot{\delta}^c - \left[\frac{3}{2} + 12\frac{p}{\rho} - \frac{9}{2}\frac{p^2}{\rho^2} - 9\frac{\dot{p}}{\dot{\rho}}\right]H^2\delta^c - \frac{1}{a^2}\frac{\Delta\hat{p}^c}{\rho} = 0\ . \tag{73}$$

Changing to a as independent variable ($\delta' \equiv \frac{d\delta^c}{da}$) and transforming into the k space, we arrive at

$$\delta^{c\prime\prime} + \left[\frac{3}{2} - \frac{15}{2}\frac{p}{\rho} + 3\frac{\dot{p}}{\dot{\rho}}\right]\frac{\delta^{c\prime}}{a} - \left[\frac{3}{2} + 12\frac{p}{\rho} - \frac{9}{2}\frac{p^2}{\rho^2} - 9\frac{\dot{p}}{\dot{\rho}}\right]\frac{\delta^c}{a^2} + \frac{k^2}{a^2H^2}\frac{\hat{p}^c}{\rho a^2} = 0\ . \tag{74}$$

Equation (74) governs the behavior of the total energy-density perturbations. As we shall see, via the $\hat{p}^c$ term the perturbations δ^c are coupled to the relative perturbations $\delta_m - D_H$.

VII. COMBINING THE SEPARATE CONSERVATION EQUATIONS

Now we combine the separate energy conservation equations (61) and (65) of the components and define $S_{mH} \equiv \delta_m - D_H$. Then

$$\dot{S}_{mH} = \Theta\left(P_H - \frac{\dot{p}_H}{\dot{\rho}_H}D_H\right) + \hat{\Theta}_H - \hat{\Theta}_m\ , \tag{75}$$

where $P_H = \frac{\hat{p}_H}{\rho_H + p_H}$. Combining the momentum balances (62) and (66) results in

$$\left(v_H - v_m\right)^{\cdot} = -P^{c_H}\ . \tag{76}$$

Because of the structure of the first-order expressions in (53) one has

$$\hat{\Theta}_H - \hat{\Theta}_m = \frac{1}{a^2}\Delta\left(v_H - v_m\right)\ . \tag{77}$$

Eq. (75) then becomes

$$\dot{S}_{mH} = \Theta\left(P_H - \frac{\dot{p}_H}{\dot{\rho}_H}D_H\right) + \frac{1}{a^2}\Delta\left(v_H - v_m\right). \tag{78}$$

Differentiation of (78) yields

$$\ddot{S}_{mH} = \left[\Theta\left(P_H - \frac{\dot{p}_H}{\dot{\rho}_H}D_H\right)\right]^{\cdot} - 2H\frac{1}{a^2}\Delta\left(v_H - v_m\right) + \frac{1}{a^2}\Delta\left(v_H - v_m\right)^{\cdot} \ . \tag{79}$$

Using here (78) again and also (76) results in

$$\ddot{S}_{mH} + 2H\dot{S}_{mH} = \left[\Theta\left(P_H - \frac{\dot{p}_H}{\dot{\rho}_H}D_H\right)\right]^{\cdot} + 2H\Theta\left(P_H - \frac{\dot{p}_H}{\dot{\rho}_H}D_H\right) - \frac{\Delta P^{c_H}}{a^2} \ . \tag{80}$$

The difference $P_H - \frac{\dot{p}_H}{\dot{\rho}_H}D_H$ describes the deviation of the DE pressure perturbations from being adiabatic. It is zero for purely adiabatic DE perturbations. We discuss this issue in more detail in the following section.

VIII. NON-ADIABATICITY AND FINAL SET OF EQUATIONS

Generally, the deviation from adiabaticity in a two-component system with components m (pressureless) and H is

$$\frac{\hat{p}}{\rho + p} - \frac{\dot{p}}{\dot{\rho}}\frac{\hat{\rho}}{\rho + p} = P^c - \frac{\dot{p}}{\dot{\rho}}D^c = \frac{\rho_H + p_H}{\rho + p}\left(\frac{\hat{p}_H}{\rho_H + p_H} - \frac{\dot{p}_H}{\dot{\rho}_H}\frac{\hat{\rho}_H}{\rho_H + p_H}\right) + \frac{\rho_m\left(\rho_H + p_H\right)}{\left(\rho + p\right)^2}\frac{\dot{p}_H}{\dot{\rho}_H}\left[\frac{\hat{\rho}_H}{\rho_H + p_H} - \frac{\hat{\rho}_m}{\rho_m}\right] \ . \tag{81}$$

Let us consider the combination $\hat{p}_H - \frac{\dot{p}_H}{\dot{\rho}_H}\hat{\rho}_H$. With $p_H = \omega\rho_H$ one has

$$\hat{p}_H - \frac{\dot{p}_H}{\dot{\rho}_H}\hat{\rho}_H = \hat{\omega}\rho_H + \left(\omega - \frac{\dot{p}_H}{\dot{\rho}_H}\right)\hat{\rho}_H \ , \tag{82}$$

where ω is given by the solution (20) and the adiabatic DE sound speed $\frac{\dot{p}_H}{\dot{\rho}_H}$ by (26). Because of (26) the combination (82) then results in

$$\hat{p}_H - \frac{\dot{p}_H}{\dot{\rho}_H}\hat{\rho}_H = \rho_H\left[\hat{\omega} + \frac{\omega}{3}rD_H\right] \ , \tag{83}$$

which is a gauge-invariant expression. In general, now an assumption for the perturbed EoS parameter $\hat{\omega}$ is necessary to proceed. We shall consider here the adiabatic case

$$\hat{p}_H = \frac{\dot{p}_H}{\dot{\rho}_H}\hat{\rho}_H \qquad \Rightarrow \qquad \hat{\omega} = -\frac{r\omega}{3}D_H \ . \tag{84}$$

This assumption of an adiabatic DE component allows us to relate the otherwise undetermined perturbation $\hat{\omega}$ of the EoS parameter to the DE energy perturbation D_H. Under this circumstances equation (80) reduces to

$$\ddot{S}_{mH} + 2H\dot{S}_{mH} = -\frac{\Delta P^{c_H}}{a^2} \ . \tag{85}$$

We emphasize that the total perturbation dynamics remains non-adiabatic due to the two-component nature of the medium. With an adiabatic DE component, the general relation (81) simplifies to

$$\frac{\hat{p}^c}{\rho+p} - \frac{\dot{p}}{\dot{\rho}}\frac{\hat{\rho}^c}{\rho+p} = -\frac{\rho_m\left(\rho_H+p_H\right)}{\left(\rho+p\right)^2}\frac{\dot{p}_H}{\dot{\rho}_H}S_{mH}\ , \tag{86}$$

or

$$\hat{p}^c = \frac{\dot{p}}{\dot{\rho}}\hat{\rho}^c - \frac{\rho_m\left(\rho_H+p_H\right)}{\left(\rho+p\right)}\frac{\dot{p}_H}{\dot{\rho}_H}S_{mH} \quad \Rightarrow \quad \hat{p}^c = \frac{\dot{p}}{\dot{\rho}}\left[\hat{\rho}^c - \rho_m S_{mH}\right]\ . \tag{87}$$

Through (87) the dynamics of the total energy-density perturbations, described by Eq. (74), is coupled to S_{mH}. Explicitly,

$$\delta^{c\prime\prime} + \left[\frac{3}{2} - \frac{15}{2}\frac{p}{\rho} + 3\frac{\dot{p}}{\dot{\rho}}\right]\frac{\delta^{c\prime}}{a} - \left[\frac{3}{2} + 12\frac{p}{\rho} - \frac{9}{2}\frac{p^2}{\rho^2} - 9\frac{\dot{p}}{\dot{\rho}} - \frac{k^2}{a^2H^2}\frac{\dot{p}}{\dot{\rho}}\right]\frac{\delta^c}{a^2} = \frac{k^2}{a^2H^2}\frac{\dot{p}}{\dot{\rho}}\frac{\rho_m}{\rho}\frac{S_{mH}}{a^2}\ . \tag{88}$$

At high redshift, for $a \ll 1$ equation (88) approaches the Einstein - de Sitter limit

$$\delta^{c\prime\prime} + \frac{3}{2a}\delta^{c\prime} - \frac{3}{2a^2}\delta^c = 0 \qquad (a \ll 1)\ . \tag{89}$$

To obtain an expression for the term on the right-hand side of Eq. (85) we write

$$\hat{p}^{c_H} = \hat{p} + \dot{p}v_H = \hat{p} + \dot{p}v + \dot{p}\left(v_H - v\right) \tag{90}$$

or

$$\hat{p}^{c_H} = \hat{p}^c + \dot{p}\left(v_H - v\right)\ . \tag{91}$$

Now, for the difference $v - v_H$ we have

$$v - v_H = \frac{\rho_m}{\rho+p}\left(v_m - v_H\right)\ . \tag{92}$$

From relation (78) it follows that (in k space)

$$v_m - v_H = \frac{a^2}{k^2}\dot{S}_{mH}\ . \tag{93}$$

Then $\hat{p}^{c_H}$ is written

$$\hat{p}^{c_H} = \hat{p}^c - \dot{p}\frac{\rho_m}{\rho+p}\frac{a^2}{k^2}\dot{S}_{mH} \quad \Rightarrow \quad \hat{p}^{c_H} = \hat{p}^c + 3H\frac{\dot{p}}{\dot{\rho}}\rho_m\frac{a^2}{k^2}\dot{S}_{mH}\ . \tag{94}$$

Here we introduce $\hat{p}^c$ from (87) to obtain

$$\hat{p}^{c_H} = \frac{\dot{p}}{\dot{\rho}}\left[\rho\delta^c - \rho_m\left(S_{mH} - 3H\frac{a^2}{k^2}\dot{S}_{mH}\right)\right]\ . \tag{95}$$

By relation (95) the dynamics of S_{mH} in Eq. (80) is coupled to the dynamics of δ^c. Explicitly,

$$S''_{mH} + \left[\frac{3}{2} - 3\frac{r}{1+\omega}\frac{\dot{p}}{\dot{\rho}} - \frac{3}{2}\frac{p}{\rho}\right]\frac{S'_{mH}}{a} + \frac{r}{1+\omega}\frac{\dot{p}}{\dot{\rho}}\frac{k^2}{a^2H^2}\frac{S_{mH}}{a^2} = \left(1 + \frac{r}{1+\omega}\right)\frac{\dot{p}}{\dot{\rho}}\frac{k^2}{a^2H^2}\frac{\delta^c}{a^2} \ . \tag{96}$$

For $a \ll 1$ this equation approaches

$$S''_{mH} + \frac{3}{2a}S'_{mH} = 0 \qquad (a \ll 1) \ . \tag{97}$$

There is one decaying solution and a solution S_{mH} = const. At high redshift, according to (22), the DE behaves as dust as well and we may use approximately adiabatic initial conditions $S_{mH} \approx 0$ for the coupled system (88) and (96).

Our interest is the matter-energy perturbation. To this purpose we decompose the total energy-density perturbation δ^c according to

$$\delta^c = \frac{\rho_m}{\rho}\delta^c_m + \frac{\rho_H}{\rho}\delta^c_H \ . \tag{98}$$

Combination with $S_{mH} = \delta_m - \frac{\delta_H}{1+\omega}$ leads to

$$\delta^c_m = \frac{1}{1+\frac{\omega}{1+r}}\left[\delta^c + \frac{1+\omega}{1+r}S_{mH}\right] \ , \tag{99}$$

which describes the matter-energy perturbations as a combination of δ^c and S_{mH}. To obtain its dynamics one has to solve the coupled system of equations (88) and (96).

The result for the matter perturbations of our model can be compared with the behavior of matter perturbations in the ΛCDM model. The latter can be obtained as a limiting case from Eq. (88) for the total energy-density perturbations. Using there $p = p_\Lambda = -\rho_\Lambda$ = constant and

$$\rho = \rho_M + \rho_\Lambda \ , \qquad \frac{p}{\rho} = -\frac{\rho_\Lambda}{\rho_\Lambda + \rho_M} = -\frac{1}{1+r} \ , \qquad r = r_0 a^{-3} \qquad (\Lambda\text{CDM}) \ , \tag{100}$$

Eq. (88) reduces to

$$\delta^{c\prime\prime} + \left[\frac{3}{2} - \frac{15}{2}\frac{p}{\rho}\right]\frac{\delta^{c\prime}}{a} - \left[\frac{3}{2} + 12\frac{p}{\rho} - \frac{9}{2}\frac{p^2}{\rho^2}\right]\frac{\delta^c}{a^2} = 0 \qquad (\Lambda\text{CDM}) \ . \tag{101}$$

For the perturbations we have

$$\hat{\rho}^c = \hat{\rho}^c_m \quad \Rightarrow \quad \delta^c = \frac{\hat{\rho}^c}{\rho} = \frac{\hat{\rho}^c_m}{\rho_\Lambda + \rho_m} = \delta^c_m \frac{r}{1+r} \qquad (\Lambda\text{CDM}) \ . \tag{102}$$

Then, the dynamics of δ^c_m is governed by

$$\delta^{c\prime\prime}_m + \frac{3}{2}\frac{2+r}{1+r}\frac{\delta^{c\prime}_m}{a} - \frac{3}{2}\frac{r}{1+r}\frac{\delta^c_m}{a^2} = 0 \ , \qquad (\Lambda\text{CDM}) \ . \tag{103}$$

While for $r \ll 1$ at high redshift the behavior of δ_m^c for our Ricci-type DE model is indistinguishable from that of the ΛCDM model, the future evolution is different. The "growing" mode of the latter approaches $\delta_m^c =$ constant for $r \to 0$, independently of the scale, whereas for Ricci-DE the behavior of the corresponding quantity depends on the scale and decays and/or oscillates. It remains open, whether or not these (unwanted) oscillations will disappear if intrinsic non-adiabatic DE perturbations are included. This would amount to a modification of relation (84).

A final comment concerning the interpretation of the perturbation variable is in order here. The matter density perturbation δ_m^c in relation (99) is defined with respect to the *total* comoving gauge. To obtain the matter density perturbation, comoving with the matter velocity, $\delta_m^{c_m} = \delta_m + \frac{\dot{\rho}_m}{\rho_m} v_m$, we have to consider

$$\delta_m^c = \delta_m + \frac{\dot{\rho}_m}{\rho_m} v = \delta_m + \frac{\dot{\rho}_m}{\rho_m} v_m + \frac{\dot{\rho}_m}{\rho_m} \left(v - v_m\right) = \delta_m^{c_m} + \frac{\dot{\rho}_m}{\rho_m} \left(v - v_m\right) \ . \tag{104}$$

Since

$$v - v_m = \frac{\rho_H + p_H}{\rho + p} \left(v_H - v_m\right) \ , \qquad v_H - v_m = -\frac{a^2}{k^2} \dot{S}_{mH} \ , \tag{105}$$

the quantity of interest is

$$\delta_m^{c_m} = \frac{1}{1 + \frac{\omega}{1+r}} \left[\delta^c + \frac{1+\omega}{1+r} S_{mH}\right] - \frac{3}{1 + \frac{r}{1+\omega}} \frac{a^2 H^2}{k^2} a S'_{mH} \ . \tag{106}$$

Obviously, δ_m^c and $\delta_m^{c_m}$ differ by the last term in (106). Because of the factor $\frac{a^2H^2}{k^2}$ one expects that on scales smaller than the Hubble scale the differences between δ_m^c and $\delta_m^{c_m}$ are small.

The behavior of the quantities S_{mH}, δ^c, δ_m^c and $\delta_m^{c_m}$ together with δ_m^c of the ΛCDM model is visualized in Figures 3 and 5 for $\omega_0 = -0.9$ with $k = 0.1$ and $k = 0.01$, respectively. The parameters in Fig. 4 differ from those of Fig. 3 by a higher energy-density ratio. This indicates a weak dependence on r_0. All these perturbations show a stable behavior. Instabilities (not shown here) occur, however, for very small values of r_0 and for $\omega_0 < -1$.

All figures show a certain compensation effect between ripple-type features in S_{mH} and δ^c such that the matter perturbation curves δ_m^c and $\delta_m^{c_m}$ look rather smooth. The figures also confirm that differences between δ_m^c and $\delta_m^{c_m}$ are indeed small on the chosen scales.

In figures 3 and 4 there appear oscillations of all the perturbation quantities very close to the present time. They seem to be similar to those known from (generalized) Chaplygin-gas models. Potentially, such oscillations should be testable by weak gravitational lensing



FIG. 3: Evolution of the perturbation quantities S_{mH} (dotted line), δ^c (dash-dotted line), δ^c_m (dashed line) and $\delta^{c_m}_m$ (thin solid line) for $\omega_0 = -0.9$, $r_0 = 0.4$ and $k = 0.1$. For comparison the ΛCDM result (thick solid line) is also included.

and by their possible influence on the CMB. But they may also jeopardize the model (cf. [37, 38] for a discussion of the matter power spectrum in the context of Chaplygin-gas models). On the other hand, we have chosen here the simplest possible case, assuming the intrinsic perturbations of the DE component to be adiabatic. It might well be that this is an oversimplification and a more general model that includes intrinsic non-adiabaticities might be more realistic. Moreover, the observed matter-spectrum refers to the distribution of baryons which so far are not included in our calculation.

IX. CONCLUSIONS

By its definition, non-interacting Ricci-type DE is characterized by a necessarily time-dependent EoS parameter. This makes it an observationally testable alternative to the ΛCDM model. Moreover, it establishes a a relationship between this EoS parameter and the matter content of the Universe. Ricci-DE behaves almost as dust at high redshift. The ratio of the energy densities of DM and DE varies considerably less than for the ΛCDM model. Since the time of radiation decoupling it has changed by about one order of magnitude



FIG. 4: The same quantities as in Fig. 3 for $\omega_0 = -0.9$, $r_0 = 0.8$ and $k = 0.1$.



FIG. 5: The same quantities as in Fig. 3 for $\omega_0 = -0.9$, $r_0 = 0.4$ and $k = 0.01$.

compared with roughly nine orders of magnitude for the ΛCDM model. This amounts to a remarkable alleviation of the coincidence problem. Our statistical analysis, based on recent observational data from SNIa, BAO and $H(z)$, results in a preferred value of $c^2 \approx 0.46$

for the Ricci-DE parameter which confirms earlier studies in the literature [11]. Within a gauge-invariant analysis we calculated the matter perturbations as a combination of the total energy perturbations of the cosmic medium and the relative perturbations of the components and clarified the relation to the matter perturbations of the ΛCDM model. For physically interesting parameter choices the perturbations turned out to be stable. Depending on the scale, the results coincide with those of the ΛCDM model until redshifts of the order of $z \approx 4$ for $k = 0.1$ or $z \approx 1.5$ for $k = 0.01$. For $z < 3$ ($k = 0.1$) or for $z < 0.7$ ($k = 0.01$) the differences are substantial. On smaller scales there appear oscillations of the perturbation quantities similar to those in Chaplygin-gas cosmologies. For the moment we must leave open whether these oscillations represent a real effect inside the dark sector, potentially detectable by weak gravitational lensing and/or modifications of the CMB anisotropy spectrum, or whether they reflect a deficiency of the model, possibly due to the neglect of intrinsic non-adiabatic perturbations of the DE component.

Acknowledgments

The authors would like to thank Oliver Piattella for helpful discussions. This work was supported by the “Comisión Nacional de Ciencias y Tecnología” (Chile) through the FONDECYT Grant No. 1110230 (SdC) and No. 1090613 (RH and SdC). J.F. and W.Z. acknowledge support by “FONDECYT-Concurso incentivo a la Cooperación Internacional” No. 1090613 as well as by CNPq (Brazil) and FAPES (Brazil).

---

[1] A. G. Riess et al., Astron. J. **116**, 1009 (1998); S. J. Perlmutter et al., Astrophys. J. **517**, 565 (1999); A. G. Riess et al., Astrophys. J. **607**, 665 (2004); P. Astier et al., Astron. Astrophys. **447**, 31 (2006).

[2] G. Hinshaw et al., arXiv:1212.5226.

[3] S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989); E.J. Copeland, M. Sami, S. Tsujikawa, Int. J. Mod. Phys. D **15**, 1753 (2006).

[4] A. G. Cohen, D.B. Kaplan and A.E. Nelson, Phys. Rev. Lett. **82**, 4971 (1999).

[5] M. Li, Phys. Lett. B **603**, 1 (2004).

[6] S. D. H. Hsu, Phys. Lett. B **594**, 13 (2004).

[7] D. Pavón and W. Zimdahl, Phys. Lett. B **628**, 206 (2005).

[8] W. Zimdahl and D. Pavón, Class. Quantum Grav. **24**, 5641 (2007); W. Zimdahl, IJMPD **17**, 651 (2008).

[9] Hsien-Chung Kao, Wo-Lung Lee and Feng-Li Lin, Phys. Rev. D **71**, 123518 (2005); K. Enquist, S. Hannestad and M.S. Sloth, JCAP **0202**, (2005) 004; Yungui Gong, Bin Wang and Yuan-Zhong Zhang, Phys. Rev. D **72**, 043510 (2005); Xin Zhang and Feng-Quan Wu, Phys. Rev. D **72**, 043524 (2005); Phys. Rev. D **76**, 023502 (2007); Qiang Wu, Yungui Gong, Anzhong Wang andJ.S. Alcaniz, Phys. Lett. B **659**, 34 (2008); Shao-Feng Wu, Peng-Ming Zhang, and Guo-Hong Yang, Class. Quantum Grav. **26** (2009) 055020; Yinzhe Ma, Yan Gong and Xuelei Chen, arXiv:0901.1215.

[10] R. Brustein and G. Veneziano, Phys. Rev. Lett. **84**, 5695 (2000).

[11] C. Gao, F. Q. Wu, X. Chen and Y. G. Shen, Phys. Rev. D **79**, 043511 (2009).

[12] R-G. Cai, B. Hu and Y. Zhang, Commun. Theor. Phys. **51**, 954 (2009).

[13] Lixin Xu and Yuting Wang, JCAP **1006**, 002 (2010); arXiv:1006.0296.

[14] Xin Zhang, Phys. Rev. D**79**, 103509 (2009); arXiv:0901.2262.

[15] Rong-Jia Yang, Zong-Hong Zhu and Fengquan Wu, Int. J. Mod. Phys. A **26**, 317 (2011); arXiv:1101.4797.

[16] Chao-Jun Feng and Xin-Zhou Li, Phys. Lett. B **680**, 184 (2009); arXiv:0904.2972.

[17] K. Karwan and T. Thitapura, JCAP **1201**, 017 (2012); arXiv:1110.2451.

[18] Yuting Wang, Lixin Xu and Yuanxing Gui, Phys. Rev. D **84**, 063513 (2011); arXiv:1110.4401.

[19] Yi Zhang and Hui Li, arXiv:1003.2788.

[20] I. Durán and D. Pavón, Phys. Rev. D **83**, 023504 (2011); arXiv:1012.2986.

[21] Tian-Fu Fu, Jing-Fei Zhang, Jin-Qian Chen and Xin Zhang, Eur. Phys. J. C **72**, 1932 (2012); arXiv:1112.2350.

[22] Zhenhui Zhang, Song Li, Xiao-Dong Li, Xin Zhang and Miao Li, JCAP **1206**, 009 (2012); arXiv:1204.6135.

[23] L.P. Chimento and M.G. Richarte, Phys. Rev. D **85**, 127301 (2012); arXiv:1207.1492.

[24] L.P. Chimento, M. Forte and M.G. Richarte, arXiv:1301.2737.

[25] B. Broda, IJMPD **21**, 1250053 (2012); arXiv:1111.5785.

[26] S. del Campo, J.C. Fabris, R. Herrera and W. Zimdahl, Phys. Rev. D **83**, 123006 (2011).

[27] J. Simon, L. Verde and R. Jimenez, Phys. Rev. D **71**, 123001 (2005).

[28] D. Stern, R. Jimenez, L. Verde, M. Kamionkowski and S. A. Stanford, JCAP **1002**, 008 (2010).

[29] R. Jimenez, L. Verde, T. Treu and D. Stern, Astrophys. J. **593**, 622 (2003).

[30] T-J. Zhang, C. Ma and T. Lan, Adv. in Astron. **2010**, 184284 (2010).

[31] C. Ma and T-J. Zhang, Astrophys. J. **730**, 74 (2011).

[32] O. Farooq, D. Mania and B. Ratra, *Hubble parameter measurement constraints on dark energy*, arXiv:1211.4253.

[33] B.L. Lago, M.O. Calvão, S.E. Jorás, R.R.R. Reis, I. Waga and R. Giostri, *Type Ia supernova parameter estimation: a comparison of two approaches using current datasets*, arXiv:1104.2874.

[34] R. Amanullah, et al. Astrophys. J. **716**, 712(2010).

[35] D.L. Eisenstein et al. (SDSS) Astrophys. J. **633**, 560 (2005).

[36] C. Blake et all, *The WiggleZ Dark Energy Survey: mapping the distance-redshift relation with baryon acoustic oscillations*, arXiv:1108.2635.

[37] H.B. Sandvik, M. Tegmark, M. Zaldariaga and I. Waga, Phys. Rev. D **69**, 123524 (2004).

[38] W.S. Hipólito-Ricaldi, H.E.S. Velten and W. Zimdahl, Phys. Rev. D **82**, 063507 (2010).